

Applicant : Qiwang Xu et al.  
For : THE USE OF N-ACETYL-D-GLUCOSAMINE FOR PREPARING  
MEDICINES FOR UROGENITAL TRACT INFECTION'S  
TREATMENT AND PREVENTION  
Page : 9

In the Claims:

This listing of claims will replace all prior versions and listings of claims in the application:

1. (currently amended) ~~A use of~~ A method for treating a subject suffering from urogenital tract infection, the method comprising administering to the subject an effective amount of a medicament containing N-acetyl-D-glucosamine and and/or pharmaceutical acceptable salts thereof in the manufacture of a medicament for treatment of urogenital tract infection.
2. (currently amended) ~~The use-~~ method according to claim 1, wherein said medicament is a formulation in the form for topical administration.
3. (currently amended) ~~The use-~~ method according to claim 2, wherein said medicament is a formulation in a form selected from a group consisting of lotion, aqua, emulsion, cream, ointment, and suppository.
4. (currently amended) ~~The use-~~ method according to ~~any one of claims 1 to 3~~ claim 1, wherein ~~the~~ a daily dose of said medicament- the administration is 100 to 2000 mg of N-acetyl-D-glucosamine.
5. (currently amended) ~~The use-~~ method according to ~~any one of claims 1 to 3~~ claim 1, wherein ~~the~~ a concentration of N-acetyl-D-glucosamine in said medicament is 0.1 to 10% by weight.
6. (new) The method according to claim 2, wherein a daily dose of the administration is 100 to 2000 mg of N-acetyl-D-glucosamine.

Applicant : Qiwan Xu et al.  
For : THE USE OF N-ACETYL-D-GLUCOSAMINE FOR PREPARING  
MEDICINES FOR UROGENITAL TRACT INFECTION'S  
TREATMENT AND PREVENTION  
Page : 10

7. (new) The method according to claim 3, wherein a daily dose of the administration is 100 to 2000 mg of N-acetyl-D-glucosamine.

8. (new) The method according to claim 2, wherein a concentration of N-acetyl-D-glucosamine in said medicament is 0.1 to 10% by weight.

9. (new) The method according to claim 3, wherein a concentration of N-acetyl-D-glucosamine in said medicament is 0.1 to 10% by weight.